

Fertility & Midwifery Care Center



Surgery Instructions

da Vinci® Robot-Assisted Myomectomy

These instructions are intended to address the majority of questions that arise related to your surgery. If there appears to be a conflict between these instructions and those given to you by the hospital, please follow these instructions. If there is any doubt and/or additional questions, please feel free to contact our office by calling **260-222-7401**.

Before Your Surgery

During this surgery, the da Vinci robot will be used to remove one or more muscle tumors or “fibroids” from your uterus. This is accomplished through four (4) or five (5) small dime-sized incisions on your abdomen.

It is critical to properly prepare the intestinal tract before this surgery. This is done to make manipulating or positioning the intestinal track during the surgery easier. Most importantly, this bowel preparation regimen would allow for the safe repair of a bowel injury should it occur during the surgery.

On the day before your surgery you are to have only clear liquids. Examples include juice, tea, coffee, carbonated beverages, gelatin, popsicles, sport beverages and bouillon. ***Specifically, do not consume any milk, dairy or RED products.***

At 1:00 p.m. on the day before your surgery take four (4) Dulcolax® tablets. Dulcolax® may be purchased without a prescription and found at most pharmacies.

At 2:00 p.m. mix Miralax® 238 grams into 64 ounces of water or Gatorade® and drink one (1) eight (8) ounce glass of the mixture every 30 minutes. This should require approximately four (4) hours to consume the entire mixture. Please do your best to drink all of the mixture if possible. At some point during the four hours you will begin experiencing bowel movements of soft stool and likely diarrhea later. This will likely continue for several hours.

After midnight, you are to have nothing to eat or drink. You may be instructed to take any medications you regularly take with a small amount of water the morning of your surgery. If you take insulin for diabetes, please make certain you understand how your FMCC physician wants you to manage this medication on the morning of your surgery.

After Your Surgery

After surgery you will recover in the post-operative recovery room, where you will remain for approximately one hour. You will then move to the Pre-Post area, the same area where you received your pre-operative medications and began the morning. Once you are fully awake and recovered from the anesthesia you will move to a regular hospital room. Most patients feel well enough to go home the evening following surgery, although a few opt to sleep overnight in the hospital and are ready to go home the next morning. If you elect to stay overnight, one of the nurses from our office will see you the morning following your surgery and make certain you are ready to go home.

We will want to see you approximately six (6) weeks following your surgery. Please call our office to schedule these follow-up appointments if you don't have them scheduled prior to the surgery.

Take your pain medication as directed and resume taking any medications that you were taking prior to the surgery unless your FMCC physician has directed you otherwise. In general, pain medication should not be taken on an empty stomach. There is no advantage to **not** taking pain medication following surgery. In fact, pain often slows the recovery process, so please take your pain medication as needed.

Following surgery you may eat the foods you normally eat, but you may find that frequent small meals are best tolerated during the first few days following your surgery. It is particularly important to drink adequate amounts of liquids following surgery and to rest often.

You may remove any bandages the day following your surgery and shower/bathe as you desire. If covering your incision(s) with a bandage makes you more comfortable, feel free to do so. A small amount of redness and/or inflammation at the incision site(s) is very common. If you notice this, clean the site(s) daily with hydrogen peroxide. Some patients find it helpful to dry their incision site(s) with a hair dryer on the cool setting. If you have questions or concerns about your incision site(s) don't hesitate to contact the office.

The physicians of FMCC ask that you refrain from driving a vehicle for as long as you are taking narcotic pain medications.

You may restart your regular exercise routine as you feel appropriate, however, don't be surprised if your endurance is not what it was prior to your surgery. Return to exercise slowly and gradually increase the frequency and intensity based on how you feel. In general, if a given activity causes pain, refrain from that activity for a few days and try again.

There are no restrictions on bathing or swimming in pools or lakes following your surgery.

Following a myomectomy, it is very common to experience light vaginal bleeding (the same or less than a menstrual period) and/or a blood-tinged vaginal discharge during the first 7-10 days following surgery.

Unless your FMCC physician has advised you otherwise, there are no specific restrictions on lifting, climbing stairs, sexual intercourse, or other activities. Again, proceed slowly based on how you feel and, "Listen to your body..." You may return to work when you feel ready to do so. If you need a work release letter please contact our office

and we will be happy to take care of this for you.

NOTE: Following myomectomy surgery, it is usually necessary to delay pregnancy in order to allow the uterus to completely heal. Make certain you have discussed this with your partner/spouse and you have a reliable plan to avoid pregnancy for the specified period of time.

Call us at 260-222-7401 if you experience any of the following:

- Fever above 101.5° F
- Nausea or vomiting unrelated to pain medication
- Abdominal or pelvic pain that is not relieved by pain medication
- Prolonged vaginal bleeding that is heavier than your usual menstrual bleeding
- Large amounts of vaginal discharge beyond the initial 3-5 days following surgery
- Inability to urinate